

Relative m -ovoids of elliptic quadrics

Antonio Cossidente

Dipartimento di Matematica Informatica ed Economia

Università della Basilicata

Contrada Macchia Romana

I-85100 Potenza

Italy

antonio.cossidente@unibas.it

Francesco Pavese

Dipartimento di Meccanica Matematica e Management

Politecnico di Bari

Via Orabona, 4

I-70125 Bari

Italy

francesco.pavese@poliba.it

Abstract

Let $\mathcal{Q}^-(2n+1, q)$ be an elliptic quadric of $\text{PG}(2n+1, q)$. A *relative m -ovoid* of $\mathcal{Q}^-(2n+1, q)$ (with respect to a parabolic section $\mathcal{Q} := \mathcal{Q}(2n, q) \subset \mathcal{Q}^-(2n+1, q)$) is a subset $\mathcal{R}$ of points of $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$ such that every generator of $\mathcal{Q}^-(2n+1, q)$ not contained in $\mathcal{Q}$ meets $\mathcal{R}$ in precisely m points. A relative m -ovoid having the same size as its complement (in $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$) is called a *relative hemisystem*. We show that a nontrivial relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$ is necessarily a relative

hemisystem, forcing q to be even. Also, we construct an infinite family of relative hemisystems of $\mathcal{Q}^-(4n+1, q)$, $n \geq 2$, admitting $\mathrm{PSp}(2n, q^2)$ as an automorphism group. Finally, some applications are given.

Keywords: elliptic quadric, relative hemisystem, strongly regular graph

MSC 2010: 05B25, 51A50, 51E20, 05E30

1 Introduction

In 2011 Penttinen and Williford [12] introduced the notion of *relative hemisystem* of a Hermitian surface $\mathcal{H}(3, q^2)$, q even. A relative hemisystem of $\mathcal{H}(3, q^2)$ with respect to a symplectic subgeometry $\mathcal{W}(3, q) \subset \mathcal{H}(3, q^2)$ is a set $\mathcal{S}$ of generators of $\mathcal{H}(3, q^2)$ disjoint from $\mathcal{W}(3, q)$ such that each point of $\mathcal{H}(3, q^2) \setminus \mathcal{W}(3, q)$ lies on $q/2$ generators of $\mathcal{S}$. Relative hemisystems are interesting in view of their connection with primitive Q -polynomial association schemes not arising from distance regular graphs [12, Theorem 3]. Up to date, three infinite families of relative hemisystems of $\mathcal{H}(3, q^2)$ and a sporadic example are known [12], [4], [5], [6].

In a recent paper Bamberg and Lee [1] generalized the concept of a relative hemisystem to a *relative m -cover* of a generalized quadrangle. Let Γ be a generalized quadrangle with a subquadrangle Γ' . A relative m -cover of Γ is a set $\mathcal{S}$ of external lines with respect to Γ' such that each point of $\Gamma \setminus \Gamma'$ lies on m members of $\mathcal{S}$. Bamberg and Lee showed that a relative m -cover of a generalized quadrangle of order (q^2, q) with respect to a subquadrangle of order (q, q) must be a hemisystem. In particular, they were able to prove an analogue of Segre's result [13], [8, Section 19.3]. See also [7]. Indeed they showed that a nontrivial relative m -cover of $\mathcal{H}(3, q^2)$, with respect to a symplectic subgeometry $\mathcal{W}(3, q)$ is a relative hemisystem, forcing q to be even [1, Theorem 1.1, Corollary 1.2].

An m -ovoid of a finite polar space $\mathcal{P}$ is a set of points which meets every generator of $\mathcal{P}$ in precisely m points [2].

Dual of the generalized quadrangles $\mathcal{H}(3, q^2)$ and $\mathcal{W}(3, q)$ are the elliptic quadric $\mathcal{Q}^-(5, q)$ and the parabolic quadric $\mathcal{Q}(4, q)$, respectively. A relative m -

cover of $\mathcal{H}(3, q^2)$ corresponds then to a set of points $\mathcal{R}$ of $\mathcal{Q}^-(5, q) \setminus \mathcal{Q}(4, q)$ such that each generator of $\mathcal{Q}^-(5, q)$ not on $\mathcal{Q}(4, q)$ meets $\mathcal{R}$ in m points. In this way, the dual of a relative m -cover of $\mathcal{H}(3, q^2)$ can be considered an “affine” m -ovoid of $\mathcal{Q}^-(5, q)$.

In this paper we introduce the concept of relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$ with respect to a parabolic section $\mathcal{Q} := \mathcal{Q}(2n, q) \subset \mathcal{Q}^-(2n+1, q)$ as a generalization of the dual of a relative m -cover of $\mathcal{H}(3, q^2)$. A relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$ (with respect to $\mathcal{Q}$) is a subset $\mathcal{R}$ of points of $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$ such that every generator of $\mathcal{Q}^-(2n+1, q)$ not contained in $\mathcal{Q}$ meets $\mathcal{R}$ in precisely m points. A relative m -ovoid having the same size as its complement (in $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$) is called a *relative hemisystem*.

The paper is organized as follows. In Section 2 we show that a nontrivial relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$ is necessarily a relative hemisystem, forcing q to be even. This generalizes the main result of [1]. In Section 3 we construct an infinite family of relative hemisystems of $\mathcal{Q}^-(4n+1, q)$, $n \geq 2$, admitting $\text{PSp}(2n, q^2)$ as an automorphism group. In Section 4 we show that a relative hemisystem of $\mathcal{Q}^-(2n+1, q)$ gives rise to non-classical $(q^n - 1)/(q - 1)$ -ovoids of the symplectic polar space $\mathcal{W}(2n+1, q)$, generalizing [6, Proposition 3.1, Corollary 3.2] and, assuming $q = 2$ and $n \geq 3$, to a strongly regular graph with parameters $v = 2^{n-1}(2^n - 1)$, $k = (2^{n-2} - 1)(2^n + 1)$, $\lambda = 2(2^{n-1} + 1)(2^{n-3} - 1)$, $\mu = 2^{n-1}(2^{n-2} - 1)$.

Throughout the paper we will use the term n -space to refer to an n -dimensional projective space.

2 Relative m -ovoids of $\mathcal{Q}^-(2n+1, q)$

Let $\text{PG}(2n+1, q)$ be the projective space over the finite field $\text{GF}(q)$ and let $\mathcal{Q}^-(2n+1, q)$ be an elliptic quadric of $\text{PG}(2n+1, q)$. Let $\mathcal{Q} = \mathcal{Q}(2n, q)$ be a parabolic quadric embedded in $\mathcal{Q}^-(2n+1, q)$. Let $\perp$ denote the polarity of $\text{PG}(2n+1, q)$ induced by $\mathcal{Q}^-(2n+1, q)$. A *generator* of $\mathcal{Q}^-(2n+1, q)$ is a totally singular $(n-1)$ -space. A generator of $\mathcal{Q}^-(2n+1, q)$ either is contained in $\mathcal{Q}$ or

meets $\mathcal{Q}$ in an $(n-2)$ -space. Let $\mathcal{B}$ denote the set of generators of $\mathcal{Q}^-(2n+1, q)$ not contained in $\mathcal{Q}$. Then $|\mathcal{B}| = q(q^n - 1) \prod_{i=2}^n (q^i + 1)$.

Definition 2.1. A relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$ (with respect to $\mathcal{Q}$) is a subset $\mathcal{R}$ of points of $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$ such that every generator of $\mathcal{Q}^-(2n+1, q)$ not contained in $\mathcal{Q}$ meets $\mathcal{R}$ in m points.

A relative m -ovoid is said to be *nontrivial* or *proper* if $m \neq 0$ or $m \neq q^{n-1}$. A relative m -ovoid having the same size as its complement (in $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$) is called a *relative hemisystem*.

Lemma 2.2. A relative m -ovoid $\mathcal{R}$ of $\mathcal{Q}^-(2n+1, q)$ has size $mq(q^n - 1)$.

Proof. We count in two ways the pairs (P, b) , where $P \in \mathcal{R}$, $b \in \mathcal{B}$ and $P \in b$:

$$|\mathcal{R}| \prod_{i=2}^n (q^i + 1) = mq(q^n - 1) \prod_{i=2}^n (q^i + 1)$$

and hence $|\mathcal{R}| = mq(q^n - 1)$. □

Lemma 2.3. Let $\mathcal{Q}'$ be a parabolic quadric $\mathcal{Q}(2n, q)$ embedded in $\mathcal{Q}^-(2n+1, q)$ and let P be a point of $\mathcal{Q}^-(2n+1, q)$.

- a) If $\mathcal{Q} \cap \mathcal{Q}'$ is an elliptic quadric $\mathcal{Q}^-(2n-1, q)$, then $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n + 1)$.
- b) If $\mathcal{Q} \cap \mathcal{Q}'$ is a hyperbolic quadric $\mathcal{Q}^+(2n-1, q)$, then $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n - 1)$.
- c) If $\mathcal{Q} \cap \mathcal{Q}'$ is a cone, then $|\mathcal{R} \cap \mathcal{Q}'| = mq^n$.
- d) If $P \in \mathcal{R}$, then $|\mathcal{R} \cap P^\perp| = m(q^n + 1) - q^n$.
- e) If $P \notin \mathcal{Q} \cup \mathcal{R}$, then $|\mathcal{R} \cap P^\perp| = m(q^n + 1)$.
- f) If $P \in \mathcal{Q}$, then $|\mathcal{R} \cap P^\perp| = m(q^n - q)$.

Proof. In the first three cases, we count in two ways the pairs (P, b) , where $P \in \mathcal{R} \cap \mathcal{Q}'$, $b \in \mathcal{B}$ is a generator of $\mathcal{Q}'$ and $P \in b$:

a)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \prod_{i=1}^n (q^i + 1)$$

and hence $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n + 1)$.

b)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \left(\prod_{i=1}^n (q^i + 1) - 2 \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n - 1)$.

c)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \left(\prod_{i=1}^n (q^i + 1) - \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence $|\mathcal{R} \cap \mathcal{Q}'| = mq^n$.

d) We count in two ways the pairs (T, b) , where $T \in \mathcal{R} \cap (P^\perp \setminus \{P\})$, $b \in \mathcal{B}$, $P \in b$ and $T \in b$:

$$|\mathcal{R} \cap (P^\perp \setminus \{P\})| \prod_{i=2}^{n-1} (q^i + 1) = (m - 1) \prod_{i=2}^n (q^i + 1)$$

and hence $|\mathcal{R} \cap P^\perp| = m(q^n + 1) - q^n$.

In the remaining two cases, we count in two ways the pairs (T, b) , where $T \in \mathcal{R} \cap P^\perp$, $b \in \mathcal{B}$, $P \in b$ and $T \in b$:

e)

$$|\mathcal{R} \cap P^\perp| \prod_{i=2}^{n-1} (q^i + 1) = m \prod_{i=2}^n (q^i + 1)$$

and hence $|\mathcal{R} \cap P^\perp| = m(q^n + 1)$.

f)

$$|\mathcal{R} \cap P^\perp| \prod_{i=2}^{n-1} (q^i + 1) = m \left(\prod_{i=2}^n (q^i + 1) - \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence $|\mathcal{R} \cap P^\perp| = m(q^n - q)$. □

Theorem 2.4. *Let $\mathcal{R}$ be a proper relative m -ovoid of $\mathcal{Q}^-(2n+1, q)$. Then q is even and $\mathcal{R}$ is a relative hemisystem.*

Proof. Let $\mathcal{Q}^-(2n-1, q)$ be an elliptic quadric contained in $\mathcal{Q}$. Let $\mathcal{F}$ be the pencil of quadrics embedded in $\mathcal{Q}^-(2n+1, q)$ having as base locus $\mathcal{Q}^-(2n-1, q)$. Then $\mathcal{F}$ contains, apart from $\mathcal{Q}$, $q-2$ parabolic quadrics and two cones. Let P_1, P_2 be the vertices of the two cones. We distinguish three cases according as

$P_1, P_2 \in \mathcal{R}$, $P_1, P_2 \notin \mathcal{R}$, $P_1 \in \mathcal{R}, P_2 \notin \mathcal{R}$, respectively. If the first case occurs then, from Lemma 2.3 a), d), we have

$$(q-2)m(q^n+1)+2m(q^n+1)-2q^n=mq(q^n-1)$$

and hence $m = q^{n-1}$. If the second case occurs then, from Lemma 2.3 a), e), we have

$$(q-2)m(q^n+1)+2m(q^n+1)=mq(q^n-1)$$

and hence $m = 0$. If the third case occurs then, from Lemma 2.3 a), d), e), we have

$$(q-2)m(q^n+1)+2m(q^n+1)-q^n=mq(q^n-1)$$

and hence $2m = q^{n-1}$. It follows that q is even and $m = q^{n-1}/2$. $\square$

Remark 2.5. Theorem 2.4 does not hold true for hyperbolic quadrics as shown in the following example. Let $\mathcal{Q}^-(3, q) \subset \mathcal{Q} := \mathcal{Q}(4, q) \subset \mathcal{Q}^+(5, q)$, q even. There exists a pencil $\mathcal{P}$ of parabolic quadrics having $\mathcal{Q}^-(3, q)$ as base locus. A plane of $\mathcal{Q}^+(5, q)$ meets a parabolic quadric of $\mathcal{P}$ in a line having necessarily a point on $\mathcal{Q}^-(3, q)$ and hence meets each of the q sets $\mathcal{Q}(4, q) \setminus \mathcal{Q}^-(3, q)$ in q points, i.e., each of them is a “relative q -ovoid” of $\mathcal{Q}^+(5, q)$ with respect to $\mathcal{Q}$. Of course, gluing together i of such sets, a “relative iq -ovoid” is obtained, $1 \leq i \leq q$. Hence there exist “relative m -ovoids” of $\mathcal{Q}^+(5, q)$ with respect to $\mathcal{Q}$ that are not relative hemisystems.

In view of Theorem 2.4, we will assume for the remaining part of the paper that q is even.

Let G denote the stabilizer of $\mathcal{Q}^-(2n+1, q)$ in $\text{PGL}(2n+2, q)$. Let Σ be the hyperplane of $\text{PG}(2n+1, q)$ containing $\mathcal{Q}$ and let $N = \Sigma^\perp$. Note that N is the nucleus of $\mathcal{Q}$. Let τ be the involution of G fixing Σ pointwise and N linewise.

Corollary 2.6. *If $\mathcal{R}$ is a relative hemisystem of $\mathcal{Q}^-(2n+1, q)$, then $|\mathcal{R} \cap \mathcal{R}^\tau| = 0$.*

Proof. With the same notation introduced in the proof of Theorem 2.4 we only observe that the line P_1P_2 contains N . $\square$

Remark 2.7. Note that Theorem 2.4 and Corollary 2.6 generalize [1, Theorem 1.1 (a), (b)] when the generalized quadrangle is classical.

3 Relative hemisystems of $\mathcal{Q}^-(4n+1, q)$

In this Section we construct a relative hemisystem of $\mathcal{Q}^-(4n+1, q)$ admitting the group $\mathrm{PSp}(2n, q^2)$ as an automorphism group.

Let V be a $2n$ -dimensional vector space over $\mathrm{GF}(q^2)$ and let k be an alternating form with associated polar space $\mathcal{W}(2n-1, q^2)$. The vector space V can be considered as a $4n$ -dimensional vector space V' over $\mathrm{GF}(q)$ via the inclusion $\mathrm{GF}(q) \subset \mathrm{GF}(q^2)$. Composition of k with the trace map $T : z \in \mathrm{GF}(q^2) \mapsto z + z^q \in \mathrm{GF}(q)$ provides a new alternating form k' on V' and so we obtain a new polar space $\mathcal{W}(4n-1, q)$. Now each point in $\mathrm{PG}(2n-1, q^2)$ corresponds to a 1-dimensional vector space in V , which in turn corresponds to a 2-dimensional vector space in V' , that is a 1-dimensional projective space of $\mathrm{PG}(4n-1, q)$. Extending this map from points of $\mathrm{PG}(2n-1, q^2)$ to subspaces of $\mathrm{PG}(2n-1, q^2)$, we obtain an injective map from subspaces of $\mathrm{PG}(2n-1, q^2)$ to certain subspaces of $\mathrm{PG}(4n-1, q)$:

$$\phi : \mathrm{PG}(2n-1, q^2) \rightarrow \mathrm{PG}(4n-1, q).$$

The map ϕ is called the $\mathrm{GF}(q)$ -linear representation of $\mathrm{PG}(2n-1, q^2)$.

A *line spread* of a polar space $\mathcal{P}$ is a collection $\mathcal{S}$ of mutually disjoint lines of $\mathcal{P}$ such that each point of $\mathcal{P}$ is contained in an element of $\mathcal{S}$. The set $\mathcal{D} = \{\phi(P) \mid P \in \mathcal{W}(2n-1, q^2)\}$ is an example of line spread of $\mathcal{W}(4n-1, q)$. The incidence structure whose points are the elements of $\mathcal{D}$ and whose lines are the generators of $\mathcal{W}(4n-1, q)$ spanned by two distinct elements of $\mathcal{D}$, is isomorphic to $\mathcal{W}(2n-1, q^2)$. Denote by ρ the symplectic polarity of $\mathrm{PG}(2n-1, q^2)$ associated with $\mathcal{W}(2n-1, q^2)$ and by ρ' the symplectic polarity of $\mathrm{PG}(4n-1, q)$ associated with $\mathcal{W}(4n-1, q)$

Since q is even, the symplectic polar space $\mathcal{W}(4n-1, q)$ is isomorphic to the parabolic quadric $\mathcal{Q} = \mathcal{Q}(4n, q)$. This isomorphism can be realized by projection

of $\mathcal{Q}$ from its nucleus N onto a hyperplane not containing N [14]. Denote by μ the bijection between $\mathcal{W}(4n-1, q)$ and $\mathcal{Q}$. Thus, $\mu(\mathcal{D})$ is a line spread of $\mathcal{Q}$ admitting $\mathrm{PSp}(2n, q^2)$ as an automorphism group.

Let Σ be the ambient projective space of $\mathcal{Q}$ and consider Σ as a hyperplane of a $\mathrm{PG}(4n+1, q)$. Let $\mathcal{Q}^-(4n+1, q)$ be an elliptic quadric of $\mathrm{PG}(4n+1, q)$ such that $\mathcal{Q}^-(4n+1, q) \cap \Sigma = \mathcal{Q}$. In this way, if G denotes the stabilizer of $\mathcal{Q}^-(4n+1, q)$ in $\mathrm{PGL}(4n+2, q)$, by Witt's theorem, we can say that there exists a subgroup K of G , isomorphic to $\mathrm{PSp}(2n, q^2)$, that stabilizes the line spread $\mu(\mathcal{D})$ of $\mathcal{Q}$.

Let $\perp$ denote the orthogonal polarity of $\mathrm{PG}(4n+1, q)$ associated with $\mathcal{Q}^-(4n+1, q)$. Then $N = \Sigma^\perp$. Let τ be the involution of G fixing Σ pointwise and N linewise. We are interested in the action of K on points of $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$.

Proposition 3.1. *The group K has two orbits $\mathcal{O}_1, \mathcal{O}_2$ of the same size on points of $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$. Moreover, $\mathcal{O}_1^\perp = \mathcal{O}_2$.*

Proof. Let $P_1 \in \mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$, then $P_1^\perp$ meets $\mathcal{Q}$ in an elliptic quadric $\mathcal{Q}^-(4n-1, q)$. An easy counting argument shows that $\mathcal{Q}^-(4n-1, q)$ contains exactly $|\mathcal{Q}^-(2n-1, q^2)|$ lines of $\mu(\mathcal{D})$. If $\mathcal{E}$ is an elliptic quadric of $\mathrm{PG}(2n-1, q^2)$ inducing the polarity ρ , then $\phi(\mathcal{E})$ defines a set consisting of $|\mathcal{E}|$ lines of $\mathcal{D}$ that are contained in an elliptic quadric $\mathcal{E}'$ of $\mathrm{PG}(4n-1, q)$ that in turn induces the polarity ρ' . Note that $\mu(\mathcal{E}')$ is an elliptic quadric $\mathcal{Q}^-(4n-1, q)$ contained in $\mathcal{Q}$. Moreover, the lines of $\mu(\mathcal{D})$ contained in $\mu(\mathcal{E}')$ span a $\mathrm{PG}(4n-1, q)$. On the other hand the number of elliptic quadrics of $\mathrm{PG}(2n-1, q^2)$ inducing the polarity ρ equals the number of elliptic quadrics of $\mathrm{PG}(4n-1, q)$ contained in $\mathcal{Q}$, indeed

$$|\mathrm{PSp}(2n, q^2) : \mathrm{PGO}^-(2n, q^2)| = |\mathrm{PSp}(4n, q) : \mathrm{PGO}^-(4n, q)| = \frac{q^{2n}(q^{2n}-1)}{2}.$$

It follows that there exists a one-to-one correspondence between the elliptic quadrics of $\mathrm{PG}(2n-1, q^2)$ inducing the polarity ρ and the elliptic quadrics of $\mathrm{PG}(4n-1, q)$ contained in $\mathcal{Q}$. Since $\mathrm{PGO}^-(2n, q^2)$ is maximal in K [9], we have that the stabilizer of P_1 in K is isomorphic to $\mathrm{PGO}^-(2n, q^2)$ and hence from the Orbit-Stabilizer theorem $|P_1^K| = q^{2n}(q^{2n}-1)/2$.

Let ℓ be the line joining P_1 and N and let $\ell \cap \mathcal{Q}^-(4n+1, q) = \{P_1, P_2\}$. Since K fixes N , if there was an element $\alpha \in K$ such that $P_1^\alpha = P_2$, then α would fix ℓ and hence α would lie in the stabilizer of ℓ in K . On the other hand, since $P_1^\perp \cap \Sigma = \ell^\perp$, the stabilizer of ℓ in K coincides with the stabilizer of P_1 in K , which acts identically on ℓ . It follows that $P_2 \notin P_1^K$. In particular K has two orbits $\mathcal{O}_1$ and $\mathcal{O}_2$ on points of $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$, both of size $q^{2n}(q^{2n}-1)/2$, and if ℓ is a secant line to $\mathcal{Q}^-(4n-1, q)$ containing N , then $|\ell \cap \mathcal{O}_1| = |\ell \cap \mathcal{O}_2| = 1$. $\square$

From now on we assume that $n \geq 2$. We are now interested in determining the orbits of the group K on the set $\mathcal{L}$ consisting of lines of $\mathcal{Q}^-(4n+1, q)$ not contained in $\mathcal{Q}$.

Proposition 3.2. *The group K has three orbits on lines of $\mathcal{L}$.*

Proof. With the same notation introduced in Proposition 3.1, let $P_1 \in \mathcal{O}_1$ and let $P_2 = P_1^\tau$. Through P_1 there pass $|\mathcal{Q}^-(4n-1, q)|$ lines of $\mathcal{L}$ and, as already observed, $\Gamma = P_1^\perp \cap \Sigma$ meets $\mathcal{Q}^-(4n+1, q)$ in a $\mathcal{Q}^-(4n-1, q)$ that corresponds to an elliptic quadric $\mathcal{Q}^-(2n-1, q^2)$ inducing the polarity ρ . Note that $N \notin \Gamma$. Hence $\mathcal{Q}^-(4n-1, q)$ contains $|\mathcal{Q}^-(2n-1, q^2)|$ lines of $\mu(\mathcal{D})$. Let X denote the set of points covered by the lines of $\mu(\mathcal{D})$ that are contained in $\mathcal{Q}^-(4n-1, q)$.

Let ℓ be a line of $\mathcal{Q}^-(4n+1, q)$ passing through P_1 and meeting $\mathcal{Q}$ at the point P . Let ℓ_P be the unique line of $\mu(\mathcal{D})$ through P and let $\ell' = P_2P$. The plane $\pi = \langle \ell, \ell_P \rangle$ is either contained in $\mathcal{Q}^-(4n+1, q)$ or shares with $\mathcal{Q}^-(4n+1, q)$ the $2q+1$ points of $\ell \cup \ell_P$. Let $\pi' = \langle \ell_P, P_2 \rangle$

If $P \in X$, then π is contained in $\mathcal{Q}^-(4n+1, q)$ and the solid $\langle \pi, P_2 \rangle$ meets $\mathcal{Q}^-(4n+1, q)$ in the union of the two planes, π and π' . Let $L = \phi^{-1}(\mu^{-1}(\ell_P))$. Then $L \in \mathcal{Q}^-(2n-1, q^2)$. If r is a secant line to $\mathcal{Q}^-(2n-1, q^2)$ through L , then $\mu(\phi(r))$ is a set of q^2+1 lines of $\mu(\mathcal{D})$ partitioning the point set of a $\mathcal{Q}(4, q)$ contained in $\mathcal{Q}$. In particular $\ell_P \in \mu(\phi(r))$ and $\mathcal{Q}(4, q)$ meets $\mathcal{Q}^-(4n-1, q)$ in a hyperbolic quadric $\mathcal{Q}^+(3, q)$. Indeed, since a line of $\mu(\mathcal{D})$ is either contained in $\mathcal{Q}^-(4n-1, q)$ or it is tangent to it and $\mathcal{Q}(4, q)$ contains q^2+1 lines of $\mu(\mathcal{D})$, it follows that $|\mathcal{Q}(4, q) \cap \mathcal{Q}^-(4n-1, q)| = q^2 - 1 + 2(q+1) = (q+1)^2$. Let $\mathcal{E} = \mathcal{Q}^-(4n-5, q)$ be the elliptic quadric $\langle \mathcal{Q}^+(3, q) \rangle^{\perp_{|\Gamma}}$ and

let $\mathcal{Q}^+(5, q) = \langle \mathcal{E} \rangle^\perp \cap \mathcal{Q}^-(4n+1, q)$. By construction π and π' are planes of $\mathcal{Q}^+(5, q)$ and $\mathcal{Q}(4, q) \subset \mathcal{Q}^+(5, q)$. The stabilizer of $\mathcal{Q}(4, q)$ in K certainly contains a group $H \simeq \text{PSp}(2, q^2) \times 1 \leq \text{PSp}(2, q^2) \times \text{PSp}(2n-2, q^2)$ acting in its natural representation on the q^2+1 lines of $\mu(\mathcal{D})$ contained in $\mathcal{Q}(4, q)$. Since $N \in \langle \mathcal{Q}(4, q) \rangle$, we have that $\langle \mathcal{Q}(4, q) \rangle^\perp$ is contained in Σ . Also, since H fixes $\langle \mathcal{Q}(4, q) \rangle^\perp$ pointwise, it follows that H stabilizes $\mathcal{Q}^+(5, q)$. Under the Klein map, the action of H on points of $\mathcal{Q}^+(5, q)$ corresponds to the action of the stabilizer of an elliptic quadric $\mathcal{Q}^-(3, q)$ in $\text{PGL}(4, q)$ on lines of $\text{PG}(3, q)$. Then H has two orbits of size $q^2(q^2+1)/2$ on points of $\mathcal{Q}^+(5, q) \setminus \mathcal{Q}(4, q)$ corresponding to lines of $\text{PG}(3, q)$ that are secant or external to $\mathcal{Q}^-(3, q)$, respectively. Since π and π' share a line, they belong to different systems of generators of $\mathcal{Q}^+(5, q)$. We may assume, that π is a Latin plane and π' is a Greek plane. Hence the q^2 points of $\pi \setminus \mathcal{Q}(4, q)$ (resp. $\pi' \setminus \mathcal{Q}(4, q)$) correspond to q^2 secant lines (resp. external lines) to $\mathcal{Q}^-(3, q)$ and therefore they lie in the same H -orbit. Since $H \leq K$, the line ℓ contains q points of $\mathcal{O}_1$. Analogously, the line ℓ' contains q points of $\mathcal{O}_2$.

If $P \notin X$, then π shares with $\mathcal{Q}^-(4n+1, q)$ the $2q+1$ points of $\ell \cup \ell_P$. Let $L = \phi^{-1}(\mu^{-1}(\ell_P))$. Then $L \notin \mathcal{Q}^-(2n-1, q^2)$. If r is an external line to $\mathcal{Q}^-(2n-1, q^2)$ through L , then $\mu(\phi(r))$ is a set of q^2+1 lines of $\mu(\mathcal{D})$ partitioning the point set of a $\mathcal{Q}(4, q)$ contained in $\mathcal{Q}$. In particular $\ell_P \in \mu(\phi(r))$ and $\mathcal{Q}(4, q)$ meets $\mathcal{Q}^-(4n-1, q)$ in an elliptic quadric $\mathcal{Q}^-(3, q)$. Indeed, since a line of $\mu(\mathcal{D})$ is either contained in $\mathcal{Q}^-(4n-1, q)$ or it is tangent to it and $\mathcal{Q}(4, q)$ contains q^2+1 lines of $\mu(\mathcal{D})$, it follows that $|\mathcal{Q}(4, q) \cap \mathcal{Q}^-(4n-1, q)| = q^2+1$. Let $\mathcal{H} = \mathcal{Q}^+(4n-5, q)$ be the quadric $\langle \mathcal{Q}^-(3, q) \rangle^{\perp|_r}$ and let $\mathcal{Q}^-(5, q) = \langle \mathcal{H} \rangle^\perp \cap \mathcal{Q}^-(4n+1, q)$. By construction ℓ and ℓ' are lines of $\mathcal{Q}^-(5, q)$ and $\mathcal{Q}(4, q) \subset \mathcal{Q}^-(5, q)$. The stabilizer of $\mathcal{Q}(4, q)$ in K certainly contains a group $H \simeq \text{PSp}(2, q^2) \times 1 \leq \text{PSp}(2, q^2) \times \text{PSp}(2n-2, q^2)$ acting in its natural representation on the q^2+1 lines of $\mu(\mathcal{D})$ contained in $\mathcal{Q}(4, q)$. Since $N \in \langle \mathcal{Q}(4, q) \rangle$, we have that $\langle \mathcal{Q}(4, q) \rangle^\perp$ is contained in Σ . Also, since H fixes $\langle \mathcal{Q}(4, q) \rangle^\perp$ pointwise, it follows that H stabilizes $\mathcal{Q}^-(5, q)$. Under the Klein map, the action of H on points of $\mathcal{Q}^-(5, q)$ corresponds to the action of the stabilizer of an elliptic quadric $\mathcal{Q}^-(3, q)$

contained in a symplectic polar space $\mathcal{W}(3, q)$ embedded in a Hermitian surface $\mathcal{H}(3, q^2)$ in $\text{PGU}(4, q^2)$ on lines of $\mathcal{H}(3, q^2)$. From [12], H has two orbits, $\mathcal{R}_1$ and $\mathcal{R}_2$, of size $q^2(q^2 - 1)/2$ on points of $\mathcal{Q}^-(5, q) \setminus \mathcal{Q}(4, q)$ and each of them is a relative hemisystem of $\mathcal{Q}^-(5, q)$ with respect to $\mathcal{Q}(4, q)$. In particular, we may assume that $P_i \in \mathcal{R}_i$, $i = 1, 2$ and, since $H \leq K$, we have that $\mathcal{R}_i \subset \mathcal{O}_i$, $i = 1, 2$. Also, from [12], the group H acts transitively on lines of $\mathcal{Q}^-(5, q)$ that are not contained in $\mathcal{Q}(4, q)$. Therefore ℓ and ℓ' lie in the same H -orbit and both contain $q/2$ points of $\mathcal{O}_1$ and $q/2$ points of $\mathcal{O}_2$.

Let $\mathcal{L}_i$ be the set of lines of $\mathcal{L}$ containing q points of $\mathcal{O}_i$, $i = 1, 2$ and $\mathcal{L}_3$ be the set of lines of $\mathcal{L}$ containing $q/2$ points of $\mathcal{O}_1$ and $\mathcal{O}_2$. Through the point P_1 (resp. P_2), there pass $(q^{2n-2} - 1)(q^{2n} + 1)/(q - 1)$ lines of $\mathcal{L}_1$ (resp. $\mathcal{L}_2$) and $q^{2n-2}(q^{2n} + 1)$ lines of $\mathcal{L}_3$. From Proposition 3.1, the stabilizer of P_1 (resp. P_2) in K is a group J isomorphic to $\text{PGO}^-(2n, q^2)$. The group J has two orbits on points of $\mathcal{Q}^-(4n - 1, q)$, namely X and its complement. Therefore J (and hence K) permutes in a single orbit the lines of $\mathcal{L}_i$ passing through P_1 (resp. P_2), $i = 1, 2, 3$. Hence, if $\ell_i \in \mathcal{L}_i$, we have that

$$|\ell_i^K| = \frac{q^{2n}(q^{2n} - 1)(q^{2n-2} - 1)(q^{2n} + 1)}{2q(q - 1)} = \frac{q^{2n-1}(q^{2n-2} - 1)(q^{4n} - 1)}{2(q - 1)}, i = 1, 2,$$

whereas, if $\ell_3 \in \mathcal{L}_3$, we have that

$$|\ell_3^K| = \frac{2q^{2n}(q^{2n} - 1)q^{2n-2}(q^{2n} + 1)}{2q} = q^{4n-3}(q^{4n} - 1).$$

On the other hand,

$$\sum_{i=1}^3 |\ell_i^K| = |\mathcal{L}| = \frac{q^{2n-1}(q^{2n-1} - 1)(q^{4n} - 1)}{q - 1}$$

and hence $\mathcal{L}_i = \ell_i^K$ is an orbit under the action of K , $i = 1, 2, 3$. $\square$

We are ready to prove the main result of this Section.

Theorem 3.3. $\mathcal{O}_i$ is a relative hemisystem of $\mathcal{Q}^-(4n + 1, q)$, $i = 1, 2$.

Proof. Let g be a generator of $\mathcal{Q}^-(4n + 1, q)$ not contained in $\mathcal{Q}$ and let g' be the $(2n - 2)$ -space obtained by intersecting g with $\mathcal{Q}$. Let P be a point of $g \setminus g'$ and let

$\mathcal{Q}^-(4n-1, q) = P^\perp \cap \mathcal{Q}$. Then g' is a generator of $\mathcal{Q}^-(4n-1, q)$ and, as already observed, $\mathcal{Q}^-(4n-1, q)$ contains $|\mathcal{Q}^-(2n-1, q^2)|$ lines of $\mu(\mathcal{D})$. Let X denote the set of points covered by the lines of $\mu(\mathcal{D})$ that are contained in $\mathcal{Q}^-(4n-1, q)$. From [11, Theorem 1.1, Table 1], the set X is a $(q^{2n-2} - 1)/(q - 1)$ -ovoid of $\mathcal{Q}^-(4n-1, q)$, i.e., every generator of $\mathcal{Q}^-(4n-1, q)$ meets X in $(q^{2n-2} - 1)/(q - 1)$ points. If $P \in \mathcal{O}_1$, then g contains $(q^{2n-2} - 1)/(q - 1)$ lines of $\mathcal{L}_1$ passing through P and q^{2n-2} lines of $\mathcal{L}_3$ passing through P . It follows that g contains

$$\frac{(q-1)(q^{2n-2} - 1)}{q-1} + \left(\frac{q}{2} - 1\right)q^{2n-2} + 1 = \frac{q^{2n-1}}{2}$$

points of $\mathcal{O}_1$ and

$$\frac{q}{2}q^{2n-2} = \frac{q^{2n-1}}{2}$$

points of $\mathcal{O}_2$. Analogously, if $P \in \mathcal{O}_2$. □

Remark 3.4. The relative hemisystems of $\mathcal{Q}^-(4n+1, q)$ constructed in Theorem 3.3 admit $\text{PSp}(2n, q^2)$ as an automorphism group.

Remark 3.5. With the aid of Magma [3] we checked that $\mathcal{Q}^-(7, 2)$ has no relative hemisystems.

4 Applications

In this section we show that associated to a relative hemisystem of $\mathcal{Q}^-(2n+1, q)$, $n \geq 2$, is a $\left(\frac{q^n-1}{q-1}\right)$ -ovoid of $\mathcal{W}(2n+1, q)$. Also, we provide a construction of a strongly regular graph with parameters $v = 2^{n-1}(2^n-1)$, $k = (2^{n-2}-1)(2^n+1)$, $\lambda = 2(2^{n-1}+1)(2^{n-3}-1)$, $\mu = 2^{n-1}(2^{n-2}-1)$ from a relative hemisystem of $\mathcal{Q}^-(2n+1, 2)$, $n \geq 3$. Note that, when n is even, this graph has the same parameters as the strongly regular graph constructed in [10], by considering as vertices the elliptic sections of a parabolic quadric of $\text{PG}(n, 4)$.

4.1 $\left(\frac{q^n-1}{q-1}\right)$ -ovoids of $\mathcal{W}(2n+1, q)$, q even

In $\mathcal{W}(2n+1, q)$, q even, the classical example of $\left(\frac{q^n-1}{q-1}\right)$ -ovoid is an elliptic quadric $\mathcal{Q}^-(2n+1, q)$ polarizing to the symplectic polarity of $\mathcal{W}(2n+1, q)$.

Our examples of $\left(\frac{q^n-1}{q-1}\right)$ -ovoids are examples of *quasi-quadrics* in the sense that they have the same size and the same intersection numbers with respect to hyperplanes of a non-degenerate quadric of $\text{PG}(2n+1, q)$.

Let $\mathcal{Q}^-(2n+1, q)$, $n \geq 2$, be an elliptic quadric of $\text{PG}(2n+1, q)$, let $\mathcal{Q} = \mathcal{Q}(2n, q)$ be a parabolic quadric embedded in $\mathcal{Q}^-(2n+1, q)$ and let $\mathcal{R}$ be a relative hemisystem of $\mathcal{Q}^-(2n+1, q)$ with respect to $\mathcal{Q}$. Since q is even, the polarity induced by $\mathcal{Q}^-(2n+1, q)$ gives rise to a symplectic polar space $\mathcal{W}(2n+1, q)$. If δ is a non-zero element of $\text{GF}(q)$ such that $X^2 + X + \delta = 0$ is irreducible over $\text{GF}(q)$, then we may assume that $\mathcal{Q}^-(2n+1, q)$ is given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 + X_{n+1}X_{n+2} + \delta X_{n+2}^2 = 0,$$

whereas $\mathcal{Q}$ is given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 = 0.$$

Let $\mathcal{Q}^+(2n+1, q)$ be the hyperbolic quadric given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 + X_{n+1}X_{n+2} = 0.$$

Then the quadrics $\mathcal{Q}^-(2n+1, q)$ and $\mathcal{Q}^+(2n+1, q)$ share $\mathcal{Q}(2n, q)$ and generate a pencil $\mathcal{P}$ comprising $q/2$ elliptic quadrics, $q/2$ hyperbolic quadrics and a cone $\mathcal{C}$ having as basis the parabolic quadric $\mathcal{Q}(2n, q)$ and as vertex its nucleus N . Since q is even, an orthogonal polarity gives rise to a symplectic polarity. In particular, the orthogonal polarities induced by the non-degenerate quadrics of $\mathcal{P}$ in $\text{PG}(2n+1, q)$ all polarize to the same symplectic polarity $\perp$ of $\text{PG}(2n+1, q)$ which defines the symplectic polar space $\mathcal{W}(2n+1, q)$.

Each elliptic quadric $\mathcal{Q}_i^-$ of $\mathcal{P}$ is partitioned as follows: $\mathcal{Q}_i^- = \mathcal{R}_i \cup \mathcal{R}_i^r \cup \mathcal{Q}$, $i = 1 \dots q/2$, where with $\mathcal{R}_i$ and $\mathcal{R}_i^r$ we have denoted the points on $\mathcal{Q}_i^-$ corresponding to $\mathcal{R}$ and its complement in $\mathcal{Q}_i^- \setminus \mathcal{Q}$, respectively.

Let $\mathcal{Q}_i^-$ and $\mathcal{Q}_j^-$ be two elliptic quadrics of $\mathcal{P}$. Let $\mathcal{A}$ be one of the two subsets $\mathcal{R}_i$ and $\mathcal{R}_i^c$ on $\mathcal{Q}_i^-$ and let $\mathcal{B}$ be one of the two subsets $\mathcal{R}_j$ and $\mathcal{R}_j^c$ on $\mathcal{Q}_j^-$, $i \neq j$. We show that $\mathcal{X} = \mathcal{A} \cup \mathcal{B} \cup \mathcal{Q}$ is a $(q^n - 1)/(q - 1)$ -ovoid of $\mathcal{W}(2n+1, q)$. Let π be a totally isotropic n -space of $\mathcal{W}(2n+1, q)$. Three possibilities occur:

1. π meets $\mathcal{Q}$ in an $(n-1)$ -space and π is contained in $N^\perp$. There are $\prod_{i=1}^n (q^i + 1)$ such n -spaces and none of these meets $\mathcal{X}$ furtherly.
2. π meets $\mathcal{Q}$ in an $(n-1)$ -space and π is not contained in $N^\perp$. Then π will be contained in a hyperbolic quadric of $\mathcal{P}$ and π does not meet $\mathcal{X}$ furtherly.
3. π meets $\mathcal{Q}$ in an $(n-2)$ -space σ . Then $\pi \cap \mathcal{Q}_i^-$ is an $(n-1)$ -space σ_i and $\pi \cap \mathcal{Q}_j^-$ is an $(n-1)$ -space σ_j , $i \neq j$. In this case $\sigma_i \cap \sigma_j = \sigma$. Since $\mathcal{A}$ and $\mathcal{B}$ are relative hemisystems of $\mathcal{Q}_i$ and $\mathcal{Q}_j$, respectively, we have that σ_i meets $\mathcal{A}$ in $q^{n-1}/2$ points and σ_j meets $\mathcal{B}$ in $q^{n-1}/2$ points. Then, π meets $\mathcal{X}$ in $(q^n - 1)/(q - 1)$ points.

We have proved the following proposition.

Proposition 4.1. *A relative hemisystem of $\mathcal{Q}^-(2n+1, q)$ gives rise to $2^{q/2}$ non-classical $(q^n - 1)/(q - 1)$ -ovoids of $\mathcal{W}(2n+1, q)$.*

Corollary 4.2. *The set $\mathcal{X}$ is an elliptic quasi-quadric.*

Proof. The set $\mathcal{X}$ has the same size of an elliptic quadric $\mathcal{Q}^-(2n+1, q)$ of $\text{PG}(2n+1, q)$ and from [2, Theorem 11] is a two-character set with intersection numbers $h_1 = (q^n - 1)(q^n + 1)/(q - 1)$ and $h_2 = (q^n - 1)(q^n + 1)/(q - 1) - q^n$. $\square$

The above construction of non-classical $(q^n - 1)/(q - 1)$ -ovoids of $\mathcal{W}(2n+1, q)$ could be realized by taking on the two elliptic quadrics $\mathcal{Q}_i^-$ and $\mathcal{Q}_j^-$ two subsets $\mathcal{A}$ and $\mathcal{B}$ corresponding to two non-isomorphic relative hemisystems.

4.2 A strongly regular graph

Let $\mathcal{R}$ be a relative hemisystem of $\mathcal{Q}^-(2n+1, 2)$, $n \geq 3$. Here $\perp$ denotes the orthogonal polarity of $\text{PG}(2n+1, 2)$ induced by $\mathcal{Q}^-(2n+1, 2)$. Let $\mathcal{G}$ be the graph having as vertices the points of $\mathcal{R}$ with two vertices x, y being adjacent if xy is a line of $\mathcal{Q}^-(2n+1, 2)$.

Proposition 4.3. *The graph $\mathcal{G}$ is strongly regular with parameters $v = 2^{n-1}(2^n - 1)$, $k = (2^{n-2} - 1)(2^n + 1)$, $\lambda = 2(2^{n-1} + 1)(2^{n-3} - 1)$, $\mu = 2^{n-1}(2^{n-2} - 1)$.*

Proof. The number of vertices is $v = 2^{n-1}(2^n - 1)$ and from Lemma 2.3 d) we have that $k = (2^{n-2} - 1)(2^n + 1)$. Let $x, y \in \mathcal{R}$ such that x is not adjacent to y . Then $y \in (x^\tau)^\perp \setminus \{x^\tau\}$. We have that $\mu = |x^\perp \cap y^\perp \cap \mathcal{R}|$ and $x^\perp \cap y^\perp$ is an elliptic quadric $\mathcal{Q}^-(2n-1, 2)$ meeting $\mathcal{Q}$ in the cone $P\mathcal{Q}^-(2n-3, 2)$, where $P = x^\tau y \cap \mathcal{Q}$. There are three hyperplanes through $x^\perp \cap y^\perp$, namely, $x^\perp$, $y^\perp$ and $T^\perp$, where $T = \Sigma \cap xy$ and $\Sigma = \langle \mathcal{Q} \rangle$. Then $T^\perp$ meets $\mathcal{Q}$ in a cone $P\mathcal{Q}(2n-2, 2)$ containing $P\mathcal{Q}^-(2n-3, 2)$. Hence, from Lemma 2.3 c), d), $|T^\perp \cap \mathcal{R}| = 2^{2n-2}$ and $|x^\perp \cap \mathcal{R}| = |y^\perp \cap \mathcal{R}| = 2^{n-2}(2^n + 1) - 2^n$. It follows that

$$\mu + 2(2^{n-2}(2^n + 1) - 2^n - \mu) + 2^{2n-2} - \mu = 2^{n-1}(2^n - 1)$$

and hence $\mu = 2^{n-1}(2^{n-2} - 1)$. Let $x, y \in \mathcal{R}$ such that x is adjacent to y . Then $y \in x^\perp \cap \mathcal{R} \setminus \{x\}$. We have that $\lambda = |x^\perp \cap y^\perp \cap \mathcal{R} \setminus \{x, y\}|$ and $x^\perp \cap y^\perp$ is a cone $\ell\mathcal{Q}^-(2n-3, 2)$, where $\ell = xy$, meeting $\mathcal{Q}$ in the cone $T\mathcal{Q}^-(2n-3, 2)$, with $T = \ell \cap \mathcal{Q}$. There are three hyperplanes through $x^\perp \cap y^\perp$, namely, $x^\perp$, $y^\perp$ and $T^\perp$. Hence, from Lemma 2.3 d), f), $|T^\perp \cap \mathcal{R}| = 2^{n-1}(2^{n-1} - 1)$ and $|x^\perp \cap \mathcal{R}| = |y^\perp \cap \mathcal{R}| = 2^{n-2}(2^n + 1) - 2^n$. It follows that

$$\lambda + 2 + 2(2^{n-2}(2^n + 1) - 2^n - \lambda - 2) + (2^{n-1}(2^{n-1} - 1) - \lambda - 2) = 2^{n-1}(2^n - 1)$$

and hence $\lambda = 2^{2n-3} - 2^n + 2^{n-2} - 2 = 2(2^{n-1} + 1)(2^{n-3} - 1)$. $\square$

Remark 4.4. The graph $\mathcal{G}$ arising from the relative hemisystem of $\mathcal{Q}^-(9, 2)$ constructed in Theorem 3.3 has the same parameters of the graph constructed in [10] by considering the elliptic sections of a parabolic quadric of $\text{PG}(4, 4)$. With the aid of Magma [3] we checked that the two graphs have the same automorphism group isomorphic to $2 \cdot \text{PSp}(4, 4)$. Hence it is plausible that the two graphs are isomorphic.

References

- [1] J. Bamberg, M. Lee, A relative m -cover of a Hermitian surface is a relative hemisystem, *arXiv 1608.03055v1*.

- [2] J. Bamberg, S. Kelly, M. Law, T. Penttila, Tight sets and m -ovoids of finite polar spaces, *J. Combin. Theory Ser. A* 114 (7) (2007), 1293-1314.
- [3] J. Cannon, C. Playoust, *An introduction to MAGMA*, University of Sidney, Sidney, Australia, 1993.
- [4] A. Cossidente, Relative hemisystems on the Hermitian surface, *J. Algebraic Combin.* 38 (2013) 2, 275-284.
- [5] A. Cossidente, A new family of relative hemisystems on the Hermitian surface, *Des. Codes Cryptogr.* 75 (2015) 2, 213-221.
- [6] A. Cossidente, F. Pavese, Intriguing sets of $\mathcal{W}(5, q)$, q even, *J. Combin. Theory Ser. A* 127 (2014), 303-313.
- [7] A. Cossidente, T. Penttila, Hemisystems on the Hermitian surface, *J. Lond. Math. Soc.* 72 (2005), 731-741.
- [8] J. W. P. Hirschfeld, *Finite Projective Spaces of Three Dimensions*, Oxford University Press, Oxford, 1991.
- [9] R.H. Dye, On the maximality of the orthogonal groups in the symplectic groups in characteristic two, *Math. Z.* 172 (1980) 3, 203-212.
- [10] X. Hubaut, R. Metz, A class of strongly regular graphs related to orthogonal groups, *Combinatorics 1981 (Rome, 1981)*, 469-472, *Ann. Discrete Math.*, 18, North-Holland, Amsterdam-New York, 1983.
- [11] S. Kelly, Constructions of intriguing sets of polar spaces from field reduction and derivation, *Des. Codes Cryptogr.* 43 (2007) 1, 1-8.
- [12] T. Penttila, J. Williford, New families of Q -polynomial association schemes, *J. Combin. Theory Ser. A* 118 (2011), 502-509.
- [13] B. Segre, Forme e geometrie Hermitiane con particolare riguardo al caso finito, *Ann. Mat. Pura Appl.* 70 (1965), 1-201.
- [14] J.A. Todd, As it might have been, *Bull. London Math. Soc.* 2 (1970) 4, 1-4.